

# Recent progress on HQET lagrangian

A. G. Grozin

Budker Institute of Nuclear Physics, Novosibirsk 630090, Russia

HQET lagrangian up to  $1/m^3$  terms is discussed. Consequences of reparameterization invariance are considered. Results for the chromomagnetic interaction coefficient at two loops, and in all orders in the large- $\beta_1$  approximation, are presented.

## 1 HQET lagrangian

QCD problems with a single heavy quark staying approximately at rest can be conveniently treated in the heavy quark effective field theory (HQET) (see [1] for review and references). We shift the energy zero level:  $E = m + \omega$ , and consider the region where residual energies  $\omega$  and momenta  $\vec{p}$  are not large:  $\omega \sim |\vec{p}| \sim \Lambda \ll m$ . The effective field theory is constructed to reproduce QCD on-shell scattering amplitudes expanded to some order  $(\Lambda/m)^n$ . This is achieved by writing down the most general effective Lagrangian consistent with the required symmetries, and tuning the coefficients to reproduce QCD on-shell amplitudes. Terms with  $D_0 Q$  can be eliminated by field redefinitions.

The most general lagrangian up to  $1/m^3$  is [2]–[6]

$$\begin{aligned}
 L = & Q^+ i D_0 Q \\
 & + \frac{C_k}{2m} Q^+ \vec{D}^2 Q + \frac{C_m}{2m} Q^+ \vec{B} \cdot \vec{\sigma} Q + \frac{iC_s}{8m^2} Q^+ (\vec{D} \times \vec{E} - \vec{E} \times \vec{D}) \cdot \vec{\sigma} Q + \frac{C_d}{8m^2} Q^+ [\vec{D} \cdot \vec{E}] Q \\
 & + \frac{C_{k2}}{8m^3} Q^+ \vec{D}^4 Q + \frac{C_{w1}}{8m^3} Q^+ \{ \vec{D}^2, \vec{B} \cdot \vec{\sigma} \} Q - \frac{C_{w2}}{4m^3} Q^+ D^i \vec{B} \cdot \vec{\sigma} D^i Q \\
 & + \frac{C_{p'p}}{8m^3} Q^+ (\vec{D} \vec{B} \cdot \vec{D} + \vec{D} \cdot \vec{B} \vec{D}) \cdot \vec{\sigma} Q + \frac{iC_M}{8m^3} Q^+ (\vec{D} \cdot [\vec{D} \times \vec{B}] + [\vec{D} \times \vec{B}] \cdot \vec{D}) Q \\
 & + \frac{C_{a1}}{8m^3} Q^+ (\vec{B}^2 - \vec{E}^2) Q - \frac{C_{a2}}{16m^3} Q^+ \vec{E}^2 Q + \frac{C_{a3}}{8m^3} Q^+ \text{Tr}(\vec{B}^2 - \vec{E}^2) Q - \frac{C_{a4}}{16m^3} Q^+ \text{Tr} \vec{E}^2 Q \\
 & + \frac{iC_{b1}}{8m^3} Q^+ (\vec{B} \times \vec{B} - \vec{E} \times \vec{E}) \cdot \vec{\sigma} Q - \frac{iC_{b2}}{8m^3} Q^+ (\vec{E} \times \vec{E}) \cdot \vec{\sigma} Q + \dots
 \end{aligned} \tag{1}$$

where  $Q$  is 2-component heavy-quark field. Here heavy-light contact interactions are omitted, as well as operators involving only light fields.

HQET can be rewritten in relativistic notations. Momenta of all states are decomposed as  $p = mv + k$  where residual momenta  $k \sim \Lambda$ . The heavy-quark field is now Dirac spinor obeying  $\not{v} Q_v = Q_v$ . The lagrangian is

$$\begin{aligned}
 L_v = & \bar{Q}_v i v \cdot D Q_v - \frac{C_k}{2m} \bar{Q}_v D_\perp^2 Q_v - \frac{C_m}{4m} \bar{Q}_v G_{\mu\nu} \sigma^{\mu\nu} Q_v \\
 & + \frac{iC_s}{8m^2} \bar{Q}_v \{ D_\perp^\mu, G^{\lambda\nu} \} v_\lambda \sigma_{\mu\nu} Q_v - \frac{C_d}{8m^2} \bar{Q}_v v^\mu [D_\perp^\nu G_{\mu\nu}] Q_v + \dots
 \end{aligned} \tag{2}$$

where  $D_\perp = D - v(vD)$ . The velocity  $v$  may be changed by an amount  $\delta v \lesssim \Lambda/m$  without spoiling the applicability of HQET and changing its predictions. This reparameterization invariance relates coefficients of varying degrees in  $1/m$  [7]–[13].

At the tree level, there are easier ways to find the coefficients  $C_i$  than QCD/HQET matching: Foldy-Wouthuysen transformation [14, 15], or using equations of motion [5]

(or integrating out lower components [16, 17]) followed by a field redefinition. The result is

$$\begin{aligned} C_k &= C_m = C_d = C_s = C_{k2} = C_{w1} = C_{a1} = C_{b1} = 1, \\ C_{w2} &= C_{p'p} = C_M = C_{a2} = C_{a3} = C_{a4} = C_{b2} = 0. \end{aligned} \quad (3)$$

However, these algebraic methods don't generalize to higher loops.

At  $1/m$  level, the kinetic coefficient  $C_k = 1$  due to the reparameterization invariance [7]. One-loop matching for the chromomagnetic coefficient  $C_m$  was done in [3]; two-loop anomalous dimension of the chromomagnetic operator in HQET was obtained in [18, 19], and two-loop matching was done in [19]; in [20], all orders of perturbation theory for  $C_m$  were summed at large  $\beta_1$ .

At  $1/m^2$  level, the spin-orbit coefficient  $C_s = 2C_m - 1$  due to the reparameterization invariance [21]–[24]. The Darwin term reduces to a contact interaction. One-loop matching for the heavy-light contact interactions was done in [24]. The one-loop anomalous dimension matrix of dimension 6 terms in the HQET lagrangian was obtained in [15], [22]–[25].

At  $1/m^3$  level, one-loop matching was done in [6] for the terms involving the heavy-quark fields twice and the gluon field once. The one-loop renormalization of dimension 7 terms in the HQET lagrangian was recently considered [26].

## 2 Matching quark-quark vertex

Renormalized QCD on-shell quark-quark proper vertex

$$- \bar{u}(\not{p} - m)u \quad (4)$$

gets no correction in the on-shell renormalization scheme. QCD spinors are related to HQET spinors by the Foldy-Wouthuysen transformation

$$u = \left( 1 + \frac{\not{k}}{2m} + \frac{k^2}{4m^2} + \dots \right) u_v, \quad \not{k}u_v = u_v. \quad (5)$$

Expressing QCD proper vertex via HQET spinors, we obtain

$$\bar{u}_v \frac{\vec{k}^2}{2m} u_v + \dots \quad (6)$$

Let's denote the sum of bare 1-particle-irreducible self-energy diagrams of the heavy quark in HQET at  $1/m^0$  as  $-i\frac{1+\not{p}}{2}\Sigma(\omega)$ ,  $\omega = kv$ . At the  $1/m$  level, self-energy diagrams with a single chromomagnetic vertex vanish. Let the sum of bare diagrams with a single kinetic vertex be  $-i\frac{C_k}{2m}\frac{1+\not{p}}{2}\Sigma_k(\omega, k_\perp^2)$ . Consider variation of  $\Sigma$  at  $v \rightarrow v + \delta v$  for an infinitesimal  $\delta v$  ( $v\delta v = 0$ ). All factors  $\frac{1+\not{p}}{2}$  can be combined into a single one, and the variation  $\delta\phi$  in it provides the variation of the  $\gamma$ -matrix structure in front of  $\Sigma$ . There are two sources of the variation of  $\Sigma$ . Terms from the expansion of denominators of the propagators produce insertions  $ik\delta v$ . Terms from the vertices produce  $igt^a\delta v^\mu$ . Now consider variation of  $\Sigma_k$  at  $k_\perp \rightarrow k_\perp + \delta k_\perp$  for an infinitesimal  $\delta k_\perp$ . Quark-quark kinetic

vertices produce  $i\frac{C_k}{m}k\delta k_\perp$ ; quark–quark–gluon kinetic vertices produce  $i\frac{C_k}{m}gt^a\delta k_\perp^\mu$ ; two–gluon vertices produce nothing. Therefore,

$$\frac{\partial\Sigma_k}{\partial k_\perp^\mu} = 2\frac{\partial\Sigma}{\partial v^\mu}. \quad (7)$$

This is the Ward identity of the reparameterization invariance first derived in [10]. Taking into account  $\frac{\partial\Sigma_k}{\partial k_\perp^\mu} = 2\frac{\partial\Sigma_k}{\partial k_\perp^2}k_\perp^\mu$  and  $\frac{\partial\Sigma}{\partial v^\mu} = \frac{d\Sigma}{d\omega}k_\perp^\mu$ , we obtain

$$\frac{\partial\Sigma_k}{\partial k_\perp^2} = \frac{d\Sigma}{d\omega}. \quad (8)$$

The right–hand side does not depend on  $k_\perp^2$ , and hence

$$\Sigma_k(\omega, k_\perp^2) = \frac{d\Sigma(\omega)}{d\omega}k_\perp^2 + \Sigma_{k0}(\omega). \quad (9)$$

This result can also be understood in a more direct way. Only diagrams with a quark–quark kinetic vertex contain  $k_\perp^2$ ; its coefficient is  $i\frac{C_k}{2m}$ . The sum of diagrams with a unit insertion is  $-i\frac{d\Sigma}{d\omega}$ . Note that diagrams with a quark–quark–gluon kinetic vertex vanish because there is no preferred transverse direction.

On the mass shell ( $\omega = 0$ ), the renormalized HQET quark–quark proper vertex is  $\frac{C_k}{2m}Z_Q\bar{u}_v$   $[-k_\perp^2 + \Sigma_k(0, k_\perp^2)]u_v = -\frac{C_k}{2m}Z_Q\left[1 - \frac{d\Sigma}{d\omega}\right]_{\omega=0}k_\perp^2\bar{u}_vu_v$ . On the mass shell, only diagrams with finite–mass particles in loops contribute (e.g.,  $c$ –quark loops in  $b$ –quark HQET) (Fig. 1). Taking into account  $Z_Q^{-1} = 1 - \frac{d\Sigma}{d\omega}|_{\omega=0}$  and comparing with (6), we finally obtain

$$C_k(\mu) = 1. \quad (10)$$

This argument works for an arbitrary  $\mu$ ; hence, the anomalous dimension of the kinetic–energy operator in HQET vanishes exactly. In a similar way, it is not difficult to prove that

$$C_{k2} = 1. \quad (11)$$

$$\begin{aligned} Z_Q & \left[ \text{tree} + \text{gluon loop } k^2 + \text{gluon loop } k + \text{gluon loop } k + \text{gluon loop } 1 + \dots \right] = \\ & Z_Q \left[ 1 + \text{gluon loop } k^2 + \dots \right] C_k \frac{k^2}{2m} \\ & Z_Q^{-1} = 1 - \frac{d\Sigma}{d\omega} = 1 + \text{gluon loop } k^2 + \dots \end{aligned}$$

Figure 1: HQET quark–quark proper vertex on the mass shell

### 3 Matching quark–quark–gluon vertex

QCD on-shell proper vertex is characterized by 2 form factors:

$$\begin{aligned} \bar{u}(p')t^a \left( \varepsilon(q^2) \frac{(p+p')^\mu}{2m} + \mu(q^2) \frac{[\not{q}, \gamma^\mu]}{4m} \right) u(p), \\ \varepsilon(q^2) = 1 + \varepsilon' \frac{q^2}{m^2} + \dots, \quad \mu(q^2) = \mu + \mu' \frac{q^2}{m^2} + \dots \end{aligned} \quad (12)$$

The total colour charge of a quark  $\varepsilon(0) = 1$  due to the gauge invariance. Ward identities in the background field formalism [27] are shown in Fig. 2, where the large dot means convolution with the gluon incoming momentum  $q$  and colour polarization  $e^a$ , the second equalities are valid only for an infinitesimal  $q$  (or in the case of an abelian external field), and  $(t^a)^{bc} = if^{acb}$  in the adjoint representation. Therefore, the QCD proper vertex  $\Lambda_\mu^a(p, q) = \Lambda_\mu t^a$  obeys  $\Lambda_\mu^a q^\mu e^a = -\Sigma(p + qe^a t^a) + \Sigma(p)$  for infinitesimal  $q$ , or  $\Lambda_\mu(p, 0) = -\frac{\partial \Sigma(p)}{\partial p^\mu}$ . The form factor is projected out by  $\varepsilon(0) = Z_Q[1 + \frac{1}{4} \text{Tr} \Lambda_\mu v^\mu (1 + \not{v})]$ . On the mass shell,  $\frac{1}{4} \text{Tr} \frac{\partial \Sigma}{\partial p^\mu} = (1 - Z_Q^{-1})v_\mu$ , and hence  $\varepsilon(0) = 1$ .

$$\begin{aligned} \text{Diagram 1: } \text{quark line } p \rightarrow p+q \text{ with gluon loop} &= g e^a t^a \left[ \text{quark line } p+q \rightarrow p - \text{quark line } p \rightarrow p+q \right] = g \left[ \text{quark line } p+q \rightarrow p+q e^a t^a - \text{quark line } p \rightarrow p \right] \\ \text{Diagram 2: } \text{quark line } p \rightarrow p+q \text{ with gluon loop} &= g e^a (t^a)^{mn} \left[ \text{quark line } p+q \rightarrow p - \text{quark line } p \rightarrow p+q \right] = g \left[ \text{quark line } p+q \rightarrow p+q e^a t^a - \text{quark line } p \rightarrow p \right] \\ \text{Diagram 3: } \text{quark line } p \rightarrow p+q \text{ with gluon loop} &= g e^a \left[ \left( \text{quark line } p \rightarrow p+q \text{ with gluon loop} - \text{quark line } p \rightarrow p \right) (t^a)^{xl} \right. \\ &\quad \left. + \left( \text{quark line } p \rightarrow p+q \text{ with gluon loop} - \text{quark line } p \rightarrow p \right) (t^a)^{xm} + \left( \text{quark line } p \rightarrow p+q \text{ with gluon loop} - \text{quark line } p \rightarrow p \right) (t^a)^{xn} \right] \\ &= g \left[ \text{quark line } p \rightarrow p+q \text{ with gluon loop} - \text{quark line } p \rightarrow p + \text{quark line } p \rightarrow p+q \text{ with gluon loop} - \text{quark line } p \rightarrow p + \text{quark line } p \rightarrow p+q \text{ with gluon loop} - \text{quark line } p \rightarrow p \right] \end{aligned}$$

Figure 2: Ward identities in the background field formalism

Let's denote the sum of bare vertex diagrams in HQET at  $1/m^0$  as  $igt^a v^\mu \frac{1+\not{v}}{2} [1 + \Lambda(\omega, \Delta)]$ , where  $\Delta = qv = \omega' - \omega$ . The Ward identity for the static quark propagator is the same as for the ordinary one (Fig. 2). Therefore,  $\Delta e^a t^a \Lambda(\omega, \Delta) = -\Sigma(\omega + \Delta e^a t^a) + \Sigma(\omega)$  for infinitesimal  $\Delta$ , or

$$\Lambda(\omega, 0) = -\frac{d\Sigma(\omega)}{d\omega}. \quad (13)$$

It is interesting, that for an abelian external field  $\Lambda(\omega, \Delta) = -\frac{\Sigma(\omega+\Delta) - \Sigma(\omega)}{\Delta}$  exactly. The total colour charge of a static quark  $Z_Q[1 + \Lambda(0, 0)] = 1$ , as expected.

The  $1/m$  HQET bare proper vertex has the form

$$i \frac{C_k}{2m} g t^a \frac{1+\not{v}}{2} \left[ (1 + \Lambda_k)(p + p')_\perp^\mu + (\Lambda_{k0} + \Lambda_{k1} p_\perp^2 + \Lambda'_{k1} p_\perp'^2 + \Lambda_{k2} q_\perp^2) v^\mu \right]$$

$$+ i \frac{C_m}{4m} g t^a \frac{1 + \not{v}}{2} [\gamma^\mu, \not{q}] \frac{1 + \not{v}}{2} (1 + \Lambda_m), \quad (14)$$

where all  $\Lambda_i$  depend on  $\omega$ ,  $\Delta$ ;  $\Lambda'_{k1}(\omega, \Delta) = \Lambda_{k1}(\omega + \Delta, -\Delta)$ ;  $\Lambda_k(\omega, \Delta) = \Lambda_k(\omega + \Delta, -\Delta)$ , and similarly for  $\Lambda_{k0}$ ,  $\Lambda_{k2}$ . Similarly to the previous Section, we can see that variation of the leading vertex function at  $v \rightarrow v + \delta v$  coincides with that of the kinetic-energy vertex function at  $p_\perp \rightarrow p_\perp + \delta p_\perp$ , if  $\delta v = \frac{C_k}{m} \delta p_\perp$ . This requires

$$\Lambda_k(\omega, \Delta) = \Lambda(\omega, \Delta), \quad \Lambda'_{k1}(\omega, \Delta) = \frac{\partial \Lambda(\omega, \Delta)}{\partial \Delta} \quad (15)$$

(and hence  $\Lambda_{k1}(\omega, \Delta) = (\frac{\partial}{\partial \omega} - \frac{\partial}{\partial \Delta}) \Lambda(\omega, \Delta)$ ). The Ward identities of Fig. 2 result in

$$\Lambda_{k0}(\omega, 0) = -\frac{d\Sigma_{k0}(\omega)}{d\omega}, \quad \Lambda_{k2}(\omega, 0) = 0 \quad (16)$$

(in an abelian external field,  $\Lambda_{k0}(\omega, \Delta) = -\frac{\Sigma_{k0}(\omega + \Delta) - \Sigma_{k0}(\omega)}{\Delta}$ ,  $\Lambda_{k2}(\omega, \Delta) = 0$ ).

Reparameterization invariance relates the spin-orbit vertex function to the chromomagnetic one, but we shall not discuss details here.

The on-shell HQET vertex at the tree level is

$$\bar{u}_v(k') \left( v^\mu + C_k \frac{(k + k')^\mu}{2m} + C_m \frac{[\not{q}, \gamma^\mu]}{4m} + C_d \frac{q^2}{8m^2} v^\mu + C_s \frac{[\not{k}, \not{q}]}{8m^2} v^\mu + \dots \right) u_v(k). \quad (17)$$

As we have demonstrated above, there are no corrections to the first two terms. Other terms have corrections starting from two loops, if there is a finite-mass flavour (such as  $c$  in  $b$ -quark HQET). Expressing the on-shell QCD vertex via HQET spinors, we obtain

$$\begin{aligned} \bar{u}_v(k') & \left[ \varepsilon(q^2) \left( v^\mu + \frac{(k + k')^\mu}{2m} - \frac{q^2 + [\not{k}, \not{q}]}{8m^2} v^\mu + \dots \right) \right. \\ & \left. + \mu(q^2) \left( \frac{[\not{q}, \gamma^\mu]}{4m} + \frac{q^2 + [\not{k}, \not{q}]}{4m^2} v^\mu + \dots \right) \right] u_v(k). \end{aligned} \quad (18)$$

Therefore, the coefficients in the HQET lagrangian are

$$C_k = 1, \quad C_m = \mu, \quad C_d = 8\varepsilon' + 2\mu - 1, \quad C_s = 2\mu - 1. \quad (19)$$

The first one has no corrections (10). The coefficients (19) are not independent:

$$C_s = 2C_m - 1. \quad (20)$$

Probably, reparameterization-invariance Ward identities yield relations among corrections from finite-mass loops in HQET which ensure the absence of corrections to (20). However, we shall not trace details here.

Similarly, at the  $1/m^3$  level, the coefficients in the HQET lagrangian are

$$C_{w1} = 4\mu' + \frac{1}{2}\mu + \frac{1}{2}, \quad C_{w2} = 4\mu' + \frac{1}{2}\mu - \frac{1}{2}, \quad C_{p'p} = \mu - 1, \quad C_M = -4\varepsilon' - \frac{1}{2}\mu + \frac{1}{2}. \quad (21)$$

They are not independent:

$$C_{w2} = C_{w1} - 1, \quad C_{p'p} = C_m - 1, \quad C_M = \frac{1}{2}(C_m - C_d). \quad (22)$$

Calculation of  $C_a$ ,  $C_b$  requires matching amplitudes with two gluons. Calculation of contact terms requires matching amplitudes with light quarks.

## 4 Chromomagnetic interaction at two loops

As we know, the kinetic coefficient  $C_k(\mu) = 1$ , and the only coefficient in the HQET lagrangian up to  $1/m$  level which is not known exactly is the chromomagnetic coefficient  $V_m(\mu)$ . It is natural to find it from QCD/HQET matching at  $\mu \sim m$  where no large logarithms appear. Renormalization group can be used to obtain  $C_m$  at  $\mu \ll m$ :

$$C_m(\mu) = C_m(m) \exp \left( - \int_{\alpha_s(m)}^{\alpha_s(\mu)} \frac{\gamma_m(\alpha)}{2\beta(\alpha)} \frac{d\alpha}{\alpha} \right), \quad (23)$$

where  $C_m(m) = 1 + C_1 \frac{\alpha_s(m)}{4\pi} + C_2 \left( \frac{\alpha_s}{4\pi} \right)^2 + \dots$ ,  $\gamma_m = \frac{d \log Z_m}{d \log \mu} = \gamma_1 \frac{\alpha_s}{4\pi} + \gamma_2 \left( \frac{\alpha_s}{4\pi} \right)^2 + \dots$  is the anomalous dimension of the chromomagnetic operator in HQET, and the  $\beta$ -function is  $\beta = -\frac{1}{2} \frac{d \log \alpha_s}{d \log \mu} = \beta_1 \frac{\alpha_s}{4\pi} + \beta_2 \left( \frac{\alpha_s}{4\pi} \right)^2 + \dots$  (where  $\beta_1 = \frac{11}{3} C_A - \frac{4}{3} T_F n_f$ ). If  $L = \log m/\mu$  is not very large, it is better to retain all two-loop terms and neglect higher loops:

$$C_m(\mu) = 1 + (C_1 - \gamma_1 L) \frac{\alpha_s(m)}{4\pi} + [C_2 - (C_1 \gamma_1 + \gamma_2) L + \gamma_1 (\gamma_1 - \beta_1) L^2] \left( \frac{\alpha_s}{4\pi} \right)^2. \quad (24)$$

This approximation holds up to relatively large  $L$  because  $C_2$  is numerically large. If  $L$  is parametrically large, then it is better to sum leading and subleading logarithms:

$$C_m(\mu) = \left( \frac{\alpha_s(\mu)}{\alpha_s(m)} \right)^{-\frac{\gamma_1}{2\beta_1}} \left[ 1 + C_1 \frac{\alpha_s(m)}{4\pi} - \frac{\beta_1 \gamma_2 - \beta_2 \gamma_1}{2\beta_1^2} \frac{\alpha_s(\mu) - \alpha_s(m)}{4\pi} \right]. \quad (25)$$

In this case, we cannot utilize  $C_2$  without knowing  $\gamma_3$ . In general, the solution of (23) can be written as

$$C_m(\mu) = \hat{C}_m K(\mu), \quad \hat{C}_m = \alpha_s(m)^{\frac{\gamma_1}{2\beta_1}} (1 + \delta c), \quad \delta c = c_1 \frac{\alpha_s(m)}{4\pi} + c_2 \left( \frac{\alpha_s(m)}{4\pi} \right)^2 + \dots \quad (26)$$

where  $\hat{C}_m$  is scale- and scheme-independent.

As a simple application, we consider  $B$ - $B^*$  mass splitting [28, 29]<sup>1</sup>

$$m_{B^*} - m_B = \frac{2C_m(\mu)}{3m} \mu_m^2(\mu) + \frac{1}{3m^2} [C_m(\mu) \rho_{km}^3(\mu) + C_m^2(\mu) \rho_{mm}^3(\mu) - C_s(\mu) \rho_s^3(\mu)], \quad (27)$$

where  $\mu_m^2(\mu)$  and  $\rho_s^3(\mu)$  are local matrix elements of chromomagnetic interaction and spin-orbit one, while  $\rho_{km}^3(\mu)$  and  $\rho_{mm}^3(\mu)$  are kinetic-chromomagnetic and chromomagnetic-chromomagnetic bilocal matrix elements (in the later case, there are two  $\gamma$ -matrix structures, 1 and  $\sigma_{\mu\nu}$ ; the coefficient of the second one is implied here). Introducing renormalization group invariants

$$\begin{aligned} \hat{\mu}_m^2 &= K(\mu) \mu_m^2(\mu), & \hat{\rho}_{km}^3 &= K(\mu) \rho_{km}^3(\mu) + [1 - K(\mu)] \rho_s^3(\mu), \\ \hat{\rho}_{mm}^3 &= K^2(\mu) \rho_{mm}^3, & \hat{\rho}_s^3 &= \rho_s^3(\mu), \end{aligned} \quad (28)$$

we can rewrite it as

$$m_{B^*} - m_B = \frac{2\hat{C}_m}{3m} \hat{\mu}_m^2 + \frac{1}{3m^2} [\hat{C}_m (\hat{\rho}_{km}^3 - 2\hat{\rho}_s^3) + \hat{C}_m^2 \hat{\rho}_{mm}^3 + \hat{\rho}_s^3]. \quad (29)$$




Figure 3: Diagrams for the QCD proper vertex





There is no sensible limit of QCD in which  $\beta_1$  may be considered a large parameter (except, may be,  $n_f \rightarrow -\infty$ ). However, retaining only the leading  $\beta_1$  terms often gives a good approximation to exact multi-loop results [33]. This limit is believed to provide information about summability of perturbation series [34]. At the first order in  $1/\beta_1$ , multiplicative renormalization amounts to subtraction of  $1/\varepsilon^n$  terms;

$$\frac{\beta_1 g_0^2}{(4\pi)^2} = \bar{\mu}^{2\varepsilon} \frac{\beta}{1 + \beta/\varepsilon}, \quad \beta = \frac{\beta_1 \alpha_s}{4\pi} = \frac{1}{2 \log \mu / \Lambda_{\overline{\text{MS}}}}. \quad (36)$$

The perturbation series (35) can be rewritten as

$$C_m(\mu) = 1 + \frac{1}{\beta_1} \sum_{L=1}^{\infty} \frac{F(\varepsilon, L\varepsilon)}{L} \left( \frac{\beta}{\varepsilon + \beta} \right)^L - (\text{subtractions}) + O\left(\frac{1}{\beta_1^2}\right). \quad (37)$$

Knowledge of the function  $F(\varepsilon, u)$  allows one to obtain the anomalous dimension

$$\gamma_m = \frac{2\beta}{\beta_1} F(-\beta, 0) + O\left(\frac{1}{\beta_1^2}\right) \quad (38)$$

and the finite term

$$C_m(\mu) = 1 + \frac{1}{\beta_1} \int_{-\beta}^0 d\varepsilon \frac{F(\varepsilon, 0) - F(0, 0)}{\varepsilon} + \frac{1}{\beta_1} \int_0^{\infty} du e^{-u/\beta} \frac{F(0, u) - F(0, 0)}{u} + O\left(\frac{1}{\beta_1^2}\right) \quad (39)$$

(this method was used in [33]; see references in this paper). Renormalization group invariant (26) is

$$\delta c = \frac{1}{\beta_1} \int_0^{\infty} du e^{-\frac{4\pi}{\beta_1 \alpha_s} u} S(u) + O\left(\frac{1}{\beta_1^2}\right), \quad S(u) = e^{-\frac{5}{3}u} \frac{F(0, u) - F(0, 0)}{u} \Big|_{\mu=m} \quad (40)$$

(here  $\alpha_s$  is taken at  $\mu = m$  in the  $V$ -scheme,  $\exp(-\frac{4\pi}{\beta_1 \alpha_s} u) = (\frac{\Lambda_V}{m})^{-2u}$ ).




Figure 4:  $L$ -loop diagrams with the maximum number of quark loops.

The function  $F(\varepsilon, u)$  is determined by the coefficient of the highest degree of  $n_f$  in the  $L$ -loop term, which is given by the diagrams in Fig. 4. Calculating them, we obtain

$$\begin{aligned} F(\varepsilon, u) &= \left(\frac{\mu}{m}\right)^{2u} e^{\gamma\varepsilon} \frac{\Gamma(1+u)\Gamma(1-2u)}{\Gamma(3-u-\varepsilon)} D(\varepsilon)^{u/\varepsilon-1} N(\varepsilon, u) \\ D(\varepsilon) &= 6e^{\gamma\varepsilon} \Gamma(1+\varepsilon) B(2-\varepsilon, 2-\varepsilon) = 1 + \frac{5}{3}\varepsilon + \dots \\ N(\varepsilon, u) &= C_F 4u(1+u-2\varepsilon u) + C_A \frac{2-u-\varepsilon}{2(1-\varepsilon)} (2+3u-5\varepsilon-6\varepsilon u+2\varepsilon^2+4\varepsilon^2 u). \end{aligned} \quad (41)$$

This gives the anomalous dimension

$$\begin{aligned}\gamma_m &= C_A \frac{\alpha_s}{2\pi} \frac{\beta(1+2\beta)\Gamma(5+2\beta)}{24(1+\beta)\Gamma^3(2+\beta)\Gamma(1-\beta)} \\ &= C_A \frac{\alpha_s}{2\pi} \left[ 1 + \frac{13}{6} \frac{\beta_1 \alpha_s}{4\pi} - \frac{1}{2} \left( \frac{\beta_1 \alpha_s}{4\pi} \right)^2 + \dots \right].\end{aligned}\quad (42)$$

This perturbation series is convergent with the radius  $\beta_1 |\alpha_s| < 4\pi$ . The Borel image of  $\delta c$

$$S(u) = \frac{\Gamma(u)\Gamma(1-2u)}{\Gamma(3-u)} \left[ 4u(1+u)C_F + \frac{1}{2}(2-u)(2+3u)C_A \right] - e^{-\frac{5}{3}u} \frac{C_A}{u} \quad (43)$$

has infrared renormalon poles at  $u = \frac{n}{2}$ . They produce ambiguities in the sum of the perturbation series for  $\delta c$ , which are of order of the residues  $\sim (\Lambda_V/m)^n$ . The leading ambiguity ( $u = \frac{1}{2}$ ) is

$$\Delta \hat{C}_m = \left( 1 + \frac{7}{8} \frac{C_A}{C_F} \right) \frac{\Delta m}{m}, \quad (44)$$

where  $\Delta m$  is the ambiguity of the heavy-quark pole mass [35, 36].

Physical quantities, such as the mass splitting (27), are factorized into short-distance coefficients and long-distance hadronic matrix elements. In regularization schemes without a hard momentum cut-off, such as  $\overline{\text{MS}}$ , Wilson coefficients also contain large-distance contributions which produce infrared renormalon ambiguities. Likewise, hadronic matrix elements contain small-distance contributions which produce ultraviolet renormalon ambiguities. In other words, the separation into short- and long-distance contributions is ambiguous; only when they are combined to form a physical quantity, an unambiguous result is obtained. Cancellations between infrared and ultraviolet renormalon ambiguities in HQET were traced in [37].

Ultraviolet renormalon ambiguities in matrix elements  $\rho_i^3$  don't depend on external states, and may be calculated at the level of quarks and gluons (Fig. 5). Note that there is an ultraviolet renormalon ambiguity in the wave function renormalization  $\Delta Z_Q = \frac{3}{2} \frac{\Delta m}{m}$  (Fig. 5d). The result is

$$\Delta \rho_{km}^3 = -\frac{2}{3} \frac{C_A}{C_F} \mu_m^2 \Delta m, \quad \Delta \rho_{mm}^3 = -\frac{19}{12} \frac{C_A}{C_F} \mu_m^2 \Delta m, \quad \Delta \rho_s^3 = -\frac{1}{2} \frac{C_A}{C_F} \mu_m^2 \Delta m. \quad (45)$$

The sum of ultraviolet ambiguities of the  $1/m^2$  contributions to (27) cancels the infrared ambiguity of the leading term.

The requirement of cancellation of renormalon ambiguities in the mass splitting (28) for all  $m$  allows us to establish the structure of the leading infrared renormalon singularity in  $S(u)$  at  $u = \frac{1}{2}$  beyond the large  $\beta_1$  limit. The ultraviolet ambiguity of the square bracket in (28) should be equal to  $\hat{\mu}_m^2$  times

$$\Lambda_V = m e^{-\frac{2\pi}{\beta_1 \alpha_s}} \alpha_s^{-\frac{\beta_2}{2\beta_1^2}} [1 + O(\alpha_s)]. \quad (46)$$

In order to reproduce the correct fractional powers of  $\alpha_s$ ,  $S(u)$  in (40) should have the branch point at  $u = \frac{1}{2}$  instead of a pole:

$$S(u) = \frac{1}{\left(\frac{1}{2} - u\right)^{1+\beta_2/2\beta_1^2}} \left[ 2C_F K_1 - \frac{1}{3} C_A K_2 + \frac{19}{12} \frac{C_A K_3}{\left(\frac{1}{2} - u\right)^{-\gamma_1/2\beta_1}} + \frac{1}{2} \frac{C_A K_4}{\left(\frac{1}{2} - u\right)^{\gamma_1/2\beta_1}} \right], \quad (47)$$




Figure 5: Diagrams for  $\rho_i^3$ ; quark loops are inserted in all possible ways.

where omitted terms are suppressed as  $\frac{1}{2} - u$  compared to the displayed ones. Normalization constants are known in the large  $\beta_1$  limit only:  $K_i = 1 + O(1/\beta_1)$ . The large-order behaviour of the perturbation series for  $\delta c$  is

$$c_{n+1} = n! (2\beta_1)^n n^{\beta_2/2\beta_1^2} \left[ 4C_F K_1 - \frac{2}{3} C_A K_2 + \frac{19}{6} C_A K_3 n^{-\gamma_1/2\beta_1} + C_A K_4 n^{\gamma_1/2\beta_1} \right], \quad (48)$$

where omitted terms are suppressed as  $1/n$  compared to the displayed ones.

**Acknowledgements.** I am grateful to A. Czarnecki and M. Neubert for collaboration in writing [19, 20]; to S. Groote for ongoing collaboration; to C. Balzereit for discussing [10, 26]; to T. Mannel for useful discussions; to J. G. Körner for hospitality at Mainz during preparation of this talk; and to M. Beyer for organization of the workshop.

## References

- [1] M. Neubert, Phys. Reports **245** (1994) 259.
- [2] E. Eichten and B. Hill, Phys. Lett. **B234** (1990) 511.

- [3] E. Eichten and B. Hill, Phys. Lett. **B243** (1990) 427.
- [4] A. F. Falk, B. Grinstein, and M. E. Luke, Nucl. Phys. **B357** (1991) 185
- [5] C. L. Y. Lee, Preprint CALT-68-1663 (1991); revised (1997).
- [6] A. V. Manohar, Phys. Rev. **D56** (1997) 230.
- [7] M. Luke and A. V. Manohar, Phys. Lett. **B286** (1992) 348.
- [8] Y.-Q. Chen, Phys. Lett. **B317** (1993) 421.
- [9] W. Kilian and T. Ohl, Phys. Rev. **D50** (1994) 4649.
- [10] C. Balzereit, Diploma thesis, Darmstadt (1994).
- [11] M. Finkemeier, H. Georgi, and M. McIrvin, Phys. Rev. **D55** (1997) 6933.
- [12] R. Sundrum, Preprint BUHEP-97-14, hep-ph/9704256 (1997); Phys. Rev. **D57**, in print.
- [13] C. L. Y. Lee, Preprint UCSD-TH-97-24, hep-ph/9709238 (1997).
- [14] J. G. Körner and G. Thompson, Phys. Lett. **B264** (1991) 185.
- [15] S. Balk, J. G. Körner, and D. Pirjol, Nucl. Phys. **B428** (1994) 499.
- [16] T. Mannel, W. Roberts, and Z. Ryzak, Nucl. Phys. **B368** (19204) 92.
- [17] C. L. Y. Lee, Preprint UCSD-TH-97-23, hep-ph/9709237 (1997).
- [18] G. Amorós, M. Beneke, and M. Neubert, Phys. Lett. **B401** (1997) 81.
- [19] A. Czarnecki and A. G. Grozin, Phys. Lett. **B405** (1997) 142.
- [20] A. G. Grozin and M. Neubert, Preprint CERN-TH/97-102, hep-ph/9707318 (1997); Nucl. Phys. **B**, in print.
- [21] Y.-Q. Chen, Y. P. Kuang, and R. Oakes, Phys. Rev. **D52** (1995) 264.
- [22] C. Balzereit and T. Ohl, Phys. Lett. **B386** (1996) 335.
- [23] M. Finkemeier and M. McIrvin, Phys. Rev. **D55** (1997) 377.
- [24] B. Blok, J. G. Körner, D. Pirjol, and J. C. Rojas, Nucl. Phys. **B496** (1997) 358.
- [25] C. Bauer and A. V. Manohar, Preprint UCSD/PTH 97-19, UTP-97-17, hep-ph/9708306 (1997); Phys. Rev. **D57**, in print.
- [26] C. Balzereit, These Proceedings.
- [27] L. F. Abbott, Acta Phys. Polonica **13** (1982) 33.
- [28] T. Mannel, Phys. Rev. **D50** (1994) 428.

- [29] I. I. Bigi, M. A. Shifman, N. G. Uraltsev, and A. I. Vainshtein, Phys. Rev. **D52** (1995) 196.
- [30] N. Gray, D. J. Broadhurst, W. Grafe, and K. Schilcher, Zeit. Phys. **C48** (1990) 673.
- [31] D. J. Broadhurst, N. Gray, and K. Schilcher, Zeit. Phys. **C52** (1991) 111.
- [32] D. J. Broadhurst, Zeit. Phys. **C54** (1992) 599.
- [33] D. J. Broadhurst and A. G. Grozin, Phys. Rev. **D52** (1995) 4082.
- [34] A. H. Mueller, QCD 20 years later, ed. P. M. Zerwas and H. A. Kastrup, World Scientific (1993), p. 162.
- [35] M. Beneke and V. M. Braun, Nucl. Phys. **B426** (1994) 301.
- [36] I. I. Bigi, M. A. Shifman, N. G. Uraltsev, and A. I. Vainshtein, Phys. Rev. **D50** (1994) 2234.
- [37] M. Neubert and C. T. Sachrajda, Nucl. Phys. **B438** (1995) 235.